

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 401/12, 405/14, 401/14, A61K 31/505	A1	(11) International Publication Number: WO 97/43279 (43) International Publication Date: 20 November 1997 (20.11.97)
(21) International Application Number: PCT/EP97/02506 (22) International Filing Date: 2 May 1997 (02.05.97) (30) Priority Data: 96201283.7 10 May 1996 (10.05.96) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): BOSMANS, Jean-Paul, René, Marie, André [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). LOVE, Christopher, John [GB/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). VAN LOMMEN, Guy, Rosalia, Eugène [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).		(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: 2,4-DIAMINOPYRIMIDINE DERIVATES AS DOPAMINE D₄ RECEPTOR ANTAGONISTS**(57) Abstract**

The present invention concerns the compounds of formula (I), the N-oxide forms, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof, wherein Alk is C₁-alkanediyl or C₃-alkenediyl; R¹ is hydrogen or C₁-alkyl; R² and R³ each independently are hydrogen, C₁-alkyl or C₃-cycloalkyl; or R² and R³ may also be taken together with the nitrogen atom to which they are attached, thus forming a pyrrolidine, a piperidine or a perhydro azepine ring; R⁴ is hydrogen or halo; Q is aryl, aryloxy, di(aryl)methyl or heteroaryl; aryl is naphthyl or phenyl, said naphthyl and phenyl may optionally be substituted; and heteroaryl is quinoliny, isoquinoliny, pyridiny, thienyl, indolyl, 2,3-dihydro-1,4-benzodioxiny, 2,3-dihydro-benzofuranyl or benzodioxolanyl; said heteroaryls may optionally be substituted; it further relates to processes for their preparation, compositions comprising them as well as their use as a medicine; compounds of formula (I) containing a radioactive isotope; a process of marking dopamine D₄ receptor sites; and a process for imaging an organ are disclosed.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

2,4-DIAMINOPYRIMIDINE DERIVATES AS DOPAMINE D₄ RECEPTOR ANTAGONISTS

5 The present invention concerns 2,4-diaminopyrimidine derivatives; it further relates to processes for their preparation, compositions comprising them, as well as their use as a medicine. The compounds of the present invention exhibit specific dopamine D₄ receptor antagonism and may particularly be useful as antipsychotics, especially in the treatment and/or prevention of psychotic disorders such as schizophrenia. In addition, the present invention concerns compounds of formula (I) containing a radioactive isotope; a process of marking dopamine D₄ receptor sites; and a process for imaging an organ.


15 It is generally accepted knowledge that dopamine receptors are important for many biochemical functions in the animal body. For example, altered functions of these receptors not only participate in the genesis of psychosis, but also of anxiety, emesis, motoric functions, addiction, sleep, feeding, learning, memory, sexual behaviour, regulation of immunological responses and blood pressure. Since dopamine receptors control a great number of pharmacological events, some of which are thus far unknown, there is a possibility that compounds which exhibit a specific binding affinity for the D₄ receptor may exert a wide range of therapeutic effects in humans.

25 EP-A-0,379,806, published on August 1, 1990, discloses *N*-[2-[(4-piperidinyl)amino]-4-pyrimidinyl]benzamides and generically describes 2-[(4-piperidinyl)amino]-4-(mono- or di(alkyl)amino)-pyrimidine derivatives, all having therapeutic potential in neurological diseases of the peripheral and central nervous systems of animals. Further, WO 93/17017 published on September 2, 1993, generically discloses *N*-[1-(2,3-dihydro-(1,4-benzodioxin or benzofuranyl)-2-ylalkyl)-4-piperidinyl]-2,4-diaminopyrimidine derivatives showing 5-HT₁-like antagonistic activity.

30 The 2,4-diaminopyrimidine derivatives of the present invention surprisingly show a high degree of dopamine D₄ receptor binding affinity. Moreover, the present compounds have a selective affinity for the dopamine D₄ receptor over other dopamine receptors in the human body. The subject compounds also show variable affinity for other receptors such as, for example, the σ -binding site.

35 The present invention concerns compounds having the formula

-2-



the *N*-oxide forms, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof, wherein

Alk is C₁₋₆alkanediyl or C₃₋₆alkenediyl;

5 R¹ is hydrogen or C₁₋₄alkyl;

R² and R³ each independently are hydrogen, C₁₋₆alkyl or C₃₋₇cycloalkyl; or

R² and R³ may also be taken together with the nitrogen atom to which they are attached, thus forming a pyrrolidine, a piperidine or a perhydro azepine ring;

R⁴ is hydrogen or halo;

10 Q is aryl, aryloxy, di(aryl)methyl or heteroaryl;

aryl is naphthyl or phenyl, said naphthyl and phenyl may optionally be substituted with one, two or three substituents selected from halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkylcarbonyl, haloC₁₋₄alkyl, nitro, amino, cyano and phenyl; and

15 heteroaryl is quinoliny, isoquinoliny, pyridiny, thienyl, indolyl, 2,3-dihydro-1,4-benzodioxiny, 2,3-dihydro-benzofuranyl or benzodioxolanyl; said heteroaryls may optionally be substituted with one, two or three substituents selected from halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkylcarbonyl, haloC₁₋₄alkyl and phenyl.

As used in the foregoing definitions and hereinafter, halo is generic to fluoro, chloro, bromo and iodo; C₁₋₄alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, 1-methylethyl, 2-methylpropyl, 2,2-dimethylethyl and the like; C₁₋₆alkyl is meant to include C₁₋₄alkyl and the higher homologues thereof having 5 or 6 carbon atoms such as, for example, pentyl, 2-methylbutyl, hexyl, 2-methylpentyl and the like;

25 C₁₋₆alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, 1,1-methanediyl, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,5-pentanediy, 1,6-hexanediy, 1,2-propanediyl, 2,3-butanediyl and the like; C₃₋₆alkenediyl defines bivalent straight and branched chain hydrocarbon radicals containing one double bond and having from

30 3 to 6 carbon atoms such as, for example, 2-propen-1,3-diyl, 3-buten-1,4-diyl, 2-buten-1,4-diyl, 2-penten-1,5-diyl, 3-penten-1,5-diyl, 3-methyl-2-buten-1,4-diyl, 3-hexen-1,6-diyl and the like; and the carbon of said C₃₋₆alkenediyl connected to the nitrogen atom of the piperidine ring preferably is saturated; haloC₁₋₄alkyl is defined as polyhalo-

-3-

substituted C₁₋₄alkyl, in particular C₁₋₄alkyl substituted with 1 to 6 halogen atoms, more in particular difluoro- or trifluoromethyl.

5 The heteroaryl group represented by Q may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate. Thus, for example, when the heteroaryl group is benzodioxolanyl, it may be a 2-benzodioxolanyl, 4-benzodioxolanyl, 5-benzodioxolanyl, 6-benzodioxolanyl and 7-benzodioxolanyl; when it is quinoliny, it may be 2-quinoliny, 3-quinoliny, 4-quinoliny, 5-quinoliny, 6-quinoliny, 7-quinoliny and 8-quinoliny; when it is 10 2,3-dihydro-benzofuranyl, it may be 2,3-dihydro-benzofuran-2-yl, 2,3-dihydro-benzofuran-3-yl, 2,3-dihydro-benzofuran-4-yl, 2,3-dihydro-benzofuran-5-yl, 2,3-dihydro-benzofuran-6-yl and 2,3-dihydro-benzofuran-7-yl.

15 The pharmaceutically acceptable acid addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. Said salts can be obtained by treating the base form of the compounds of formula (I) with appropriate acids such as inorganic acids, for example, hydrohalic acid, e.g. hydrochloric or hydrobromic, sulfuric, nitric, phosphoric and the like acids; or organic acids, such as, for example, acetic, hydroxy- 20 acetic, propanoic, lactic, pyruvic, oxalic, malonic, succinic, malic, fumaric, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

25 The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

30 The *N*-oxide forms of the compounds of formula (I) are meant to comprise those compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

35 The term "stereochemically isomeric forms" as used hereinbefore and hereinafter defines all the possible isomeric forms in which the compounds of formula (I) may occur. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture, and in particular the racemic mixture, of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. Stereochemically isomeric forms of the

-4-

compounds of formula (I) and mixtures of such forms are obviously intended to be encompassed by formula (I).

Preferably, R¹ is C₁₋₄alkyl.

- 5 Preferably, R² is C₁₋₆alkyl or C₃₋₇cycloalkyl and R³ is hydrogen or C₁₋₆alkyl, or R² and R³ are taken together with the nitrogen atom to which they are attached, thus forming a pyrrolidine ring.

Preferably, Alk is a straight chained C₁₋₆alkanediyl or a straight chained C₃₋₆alkenediyl, more in particular, Alk is methylene, 1,3-propanediyl or 1,4-butanediyl.

- 10 Suitably, aryl is 2-naphthyl, 3-naphthyl, phenyl or mono- or disubstituted phenyl. Suitably, heteroaryl is optionally substituted 2-quinoliny, 3-pyridyl, 4-pyridyl, 2-thienyl, 3-indolyl, 2,3-dihydro-1,4-benzodioxin-2-yl, 2,3-dihydrobenzofuran-5-yl or 5-benzodioxolanyl.

- 15 A first group of particular compounds are those compounds of formula (I) wherein Alk is a straight chained C₁₋₆alkanediyl and Q is optionally substituted phenyl, 2-naphthyl, 2-quinoliny, 2,3-dihydrobenzofuran-5-yl or 5-benzodioxolanyl.

- 20 A second group of particular compounds are those compounds of formula (I) wherein R¹ is C₁₋₄alkyl; R² is C₁₋₆alkyl or C₃₋₇cycloalkyl and R³ is hydrogen, C₁₋₆alkyl; or R² and R³ form a pyrrolidine ring when taken together with the nitrogen atom to which they are attached.

- 25 Interesting compounds are those compounds of formula (I) wherein R⁴ is hydrogen.

- Preferred compounds are those particular compounds wherein R¹ is methyl; R² is methyl, ethyl, propyl or cyclopropyl and R³ is hydrogen, methyl or ethyl; or R² and R³ form a pyrrolidine ring when taken together with the nitrogen atom to which they are attached; Alk is methylene, and Q is optionally substituted phenyl, 2-naphthyl, 2-quinoliny, 2,3-dihydrobenzofuran-5-yl or 5-benzodioxolanyl.

- More preferred compounds are those preferred compounds wherein R² is methyl, ethyl or cyclopropyl and R³ is hydrogen or methyl; or R² and R³ form a pyrrolidine ring when taken together with the nitrogen atom to which they are attached.

35


Most preferred compounds are

N⁴-cyclopropyl-N²-methyl-N²-[1-(phenylmethyl)-4-piperidiny]-2,4-pyrimidine-diamine;

-5-

- N*²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-*N*⁴-ethyl-*N*²-methyl-2,4-pyrimidinediamine;
*N*²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-*N*⁴-cyclopropyl-*N*²-methyl-2,4-pyrimidinediamine;
 5 *N*²-[1-[(4-chlorophenyl)methyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidine diamine;
*N*²-[1-[(3-fluorophenyl)methyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidine-diamine;
 10 *N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-[[4-(trifluoromethyl)phenyl]methyl]-4-piperidiny]-2,4-pyrimidinediamine;
*N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-[(4-methylphenyl)methyl]-4-piperidiny]-2,4-pyrimidine-diamine;
*N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-[(3-methylphenyl)methyl]-4-piperidiny]-2,4-pyrimidine-diamine; the stereochemically isomeric forms and the pharmaceutically acceptable
 15 addition salts thereof.

Compounds of formula (I) can generally be prepared by *N*-alkylating an intermediate of formula (III) with an intermediate of formula (II), wherein W represents an appropriate reactive leaving group such as, for example, a halogen. The reaction can be performed
 20 in a reaction-inert solvent such as, for example, methyl isobutylketone, *N,N*-dimethylacetamide or *N,N*-dimethylformamide, in the presence of a suitable base such as, for example, sodium carbonate or triethylamine, and optionally in the presence of potassium iodide. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried out at a temperature ranging between room temperature and
 25 reflux temperature.



Compounds of formula (I) wherein Alk is attached to the nitrogen of the piperidine ring by a carbon atom bearing at least one hydrogen atom, said compounds being represented by formula (I-a) and said Alk being represented by Alk'H, can be prepared
 30 by reductively *N*-alkylating an intermediate of formula (III) with an aldehyde or ketone of formula Q-Alk'=O (IV), said Q-Alk'=O being derived from Q-Alk'H₂ by replacing two geminal hydrogen atoms by an oxo group.

[illegible]

15

$$\begin{array}{c}
 \text{Q-Alk-N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{N} \begin{array}{c} \text{R}^1 \\ | \\ \text{H} \end{array} + \text{W}^1 \begin{array}{c} \text{R}^2 \\ | \\ \text{N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{R}^3 \\ | \\ \text{R}^4 \end{array} \longrightarrow \text{Q-Alk-N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{N} \begin{array}{c} \text{R}^1 \\ | \\ \text{N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{R}^3 \\ | \\ \text{R}^4 \end{array} \\
 \text{(V)} \qquad \qquad \text{(VI)} \qquad \qquad \qquad \text{(I)}
 \end{array}$$

20 The compounds of formula (I) may be also converted into other compounds of formula (I) following art-known transformation reactions. For instance, compounds of formula (I) wherein R^4 is hydrogen, said compounds being represented by formula (I-b-1), may be converted into compounds of formula (I) wherein R^4 is a halogen atom, said compounds being represented by formula (I-b-2). For instance, an iodine atom may be


25 introduced in the 5-position of the pyrimidine ring of a compound of formula (I-b-1) by reacting said compound with a mixture of benzyltrimethylammoniumchloride and iodinechloride, or a functional analogue thereof. Alternatively, the introduction of an iodine atom in the 5-position of the pyrimidine ring of a compound of formula (I-b-1) may be established by reacting said compound with a mixture of a peroxide and I_2 .

The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

The starting materials and some of the intermediates are known compounds and are commercially available or may be prepared according to conventional reaction procedures generally known in the art. For instance, intermediates of formula (III) may be prepared according to the procedures described in WO-93/17017.


In the following paragraphs there are described several methods of preparing the intermediates employed in the foregoing preparations.

Intermediates of formula (III) can be prepared by reacting an intermediate of formula (VII), wherein P¹ is a protective group such as, e.g. alkyloxycarbonyl with an intermediate of formula (VI), following art-known *N*-alkylation methods, and subsequently removing P¹.



Intermediates of formula (V) can be prepared by reacting an intermediate of formula (VIII), wherein P² is a suitable protecting group such as, e.g. alkyloxycarbonyl, with an intermediate of formula (II), following art-known *N*-alkylation methods, and subsequently removing P².

-8-



- Some of the compounds of formula (I) and some of the intermediates in the present invention contain at least one asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers.
- 15 An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.
- Pure stereochemically isomeric forms of the compounds of formula (I) may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically. The pure and mixed stereochemically isomeric forms of the compounds of formula (I) are intended to be embraced within the scope of the present invention.
- 25 The compounds of formula (I), the *N*-oxides, the pharmaceutically acceptable acid addition salts and stereochemically isomeric forms thereof, are potent antagonists of the dopamine D₄ receptor, *i.e.* they show a high degree of dopamine D₄ receptor binding affinity thus inhibiting the binding of an endogeneous ligand, in particular dopamine, to the dopamine D₄ receptor, as is demonstrated in the pharmacological example described hereinafter. The antagonistic effect of the binding of the present compounds to the dopamine D₄ receptor was confirmed in signal-transduction assays.
- The present compounds show interesting activity in the so-called "differential reinforcement test low rate 72 seconds"-test (DRL-72) which is an *in vivo* test where most clinically active antidepressants given at high doses show activity. In said test,

rats can obtain food by pressing a lever only when they have waited a full 72 seconds between two lever presses. The present D₄ antagonists induce a more efficient behaviour of the rats whereas untreated animals find it difficult to control their impulsive tendency to press the lever and to subordinate it to appropriate timing so as to maximize their award. The usefulness of this DRL-72 test as a model for specific D₄ antagonists such as the present compounds is further supported by the fact that (a) Manki et al. (Journal of Affective Disorders 40 (1996), 7-13) found that there is a significant association between the D₄ receptor gene polymorphism and mood disorders, and (b) by the fact that D₄ receptors are known to be most dense in hippocampus, entorhinal and cerebral cortex in the primates, humans as well as rodents.

Antagonizing the dopamine D₄ receptor will suppress or relieve a variety of symptoms associated with phenomena induced by the activation, in particular the excessive activation, of said receptor. Consequently, the ability of the present compounds to alter dopamine D₄ mediated neurotransmission makes them of potential use in the treatment and/or prevention of a variety of disorders associated therewith such as sleep disorders, sexual disorders, thought disorders, impaired information processing, psychosis, affective psychosis, nonorganic psychosis, personality disorders, psychiatric mood disorders, conduct and impulse disorders, schizophrenic and schizoaffective disorders, polydipsia, bipolar disorders, dysphoric mania, anxiety and related disorders, gastrointestinal disorders, obesity, emesis, bacterial infections of the CNS such as meningitis, learning disorders, memory disorders, Parkinson's disease, depression, extrapyramidal side effects from neuroleptic agents, neuroleptic malignant syndrome, hypothalamic pituitary disorders, congestive heart failure, chemical dependencies such as drug and alcohol dependencies, vascular and cardiovascular disorders, ocular disorders, dystonia, tardive dyskinesia, Gilles De la Tourette's syndrome and other hyperkinesias, dementia, ischemia, movement disorders such as akathisia, hypertension and diseases caused by a hyperactive immune system such as allergies and inflammation.

The compounds of the present invention distinctively show affinity for the dopamine D₄ receptor in comparison with other dopamine receptors such as, for example, the dopamine D₂ receptor. Such a dissociation between dopamine D₄ receptor antagonizing activity and other dopamine receptor activity may be of additional use in the treatment and/or prevention of the above-mentioned disorders. For example, Van Tol et al. (Nature 1991, 350, 610-614) suggested that compounds which can interact selectively with the dopamine D₄ receptor, whilst having a less pronounced action at

-10-

the dopamine D₂ receptor, might have the same beneficial level of antipsychotic activity as classical antipsychotics with the additional benefit of being less prone to the undesired extrapyramidal or neuroendocrine side-effects of classical antipsychotics. It is therefore that the present compounds are particularly useful as antipsychotics, especially in the treatment and/or prevention of psychotic disorders such as schizophrenia.

In addition to their potency to antagonize the dopamine D₄ receptor, the subject compounds also show variable affinity for other receptors such as, for example, the σ -binding site.

In view of the usefulness of the subject compounds in the treatment and/or prevention of disorders associated with the excessive activation of the dopamine D₄ receptor, in particular in the treatment of psychotic disorders such as schizophrenia, the present invention provides a method of treating warm-blooded animals suffering from such diseases, said method comprising the systemic administration of a dopamine D₄ receptor antagonizing amount of a compound of formula (I), a *N*-oxide or a pharmaceutically acceptable acid addition salt thereof.

~~The present invention thus also relates to compounds of formula (I) as defined herein~~ above for use as a medicine. Further, the present invention also relates to the use of a compound of formula (I) for the manufacture of a medicament for treating psychotic disorders.

The term "dopamine D₄ receptor antagonizing amount", as used herein, refers to an amount sufficient to inhibit the binding of an endogenous ligand, in particular dopamine, to the dopamine D₄ receptor. Those of skill in the treatment of the disorders as mentioned hereinabove could determine that an effective dopamine D₄ receptor antagonizing daily amount would be from about 0.01 mg/kg to about 10 mg/kg body weight, more preferably from about 0.04 mg/kg to about 4 mg/kg body weight. The compounds may be administered on a regimen of 1 to 4 times per day.

In order to alleviate the symptoms of psychotic disorders such as schizophrenia without causing undesired side-effects, the dosage level of the compound according to the invention is ideally selected such that the dose administered is effective in substantially completely blocking the dopamine D₄ receptor while displaying a favourable dopamine D₂ receptor occupancy causing no or negligible undesired extrapyramidal or neuroendocrine side-effects.

If desired, the compounds according to this invention may be co-administered with another antipsychotic, for example one producing its effects via one or more of the following mechanisms : dopamine D₂ receptor blockade, 5-HT₂ receptor blockade, 5-HT_{1A} agonism and 5-HT₃ antagonism. In such circumstances, an enhanced anti-psychotic effect may be envisaged without a corresponding increase in side-effects such as those caused by, for example, strong dopamine D₂ receptor blockade; or a comparable antipsychotic effect with reduced side-effects may alternatively be envisaged. Such co-administration may be desirable where a patient is already established on a, for example, anti-schizophrenic treatment regime involving conventional anti-schizophrenic medicaments.

For administration purposes, the subject compounds may be formulated into various pharmaceutical forms. To prepare the pharmaceutical compositions of this invention, an effective dopamine D₄ receptor antagonizing amount of the particular compound, as acid addition salt or in its free base form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable solutions containing compounds of formula (I) may be formulated in an oil for prolonged action. Appropriate oils for this purpose are, for example, peanut oil, sesame oil, cottonseed oil, corn oil, soy bean oil, synthetic glycerol esters of long chain fatty acids and mixtures of these and other oils. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration,

-12-

the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful
5 for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment. Acid addition salts of the compounds of formula (I) due to their increased water solubility over the corresponding free base form, are more suitable in the preparation of aqueous compositions.

10

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of
15 active ingredient calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

20

Due to their high degree of specificity to the dopamine D₄ receptor, the compounds of formula (I) as defined above, are also useful to mark or identify receptors, in particular dopamine D₄ receptors. To this purpose, the compounds of the present invention need to be labelled, in particular by replacing, partially or completely, one or more atoms in
25 the molecule by their radioactive isotopes. Examples of interesting labelled compounds are those compounds having at least one halo which is a radioactive isotope of iodine, bromine or fluorine; or those compounds having at least one ¹¹C-atom or tritium atom.

One particular group consists of those compounds of formula (I) wherein Q is aryl or
30 heteroaryl substituted with a radioactive halogen atom, or those compounds wherein R⁴ is a radioactive halogen atom. In principle, any compound of formula (I) containing a halogen atom is prone for radiolabelling by replacing the halogen atom by a suitable isotope. Suitable halogen radioisotopes to this purpose are radioactive iodides, e.g. ¹²²I, ¹²³I, ¹²⁵I, ¹³¹I; radioactive bromides, e.g. ⁷⁵Br, ⁷⁶Br, ⁷⁷Br and ⁸²Br, and radioactive
35 fluorides, e.g. ¹⁸F. The introduction of a radioactive halogen atom can be performed by a suitable exchange reaction or by using any one of the procedures as described

hereinabove to prepare halogen derivatives of formula (I), in particular, the reactions for converting compounds of formula (I-b-1) to compounds of formula (I-b-2).

5 Preferred labelled compounds are those compounds of formula (I), wherein Q is aryl or heteroaryl substituted with ^{123}I , ^{125}I , ^{75}Br , ^{76}Br , ^{77}Br or ^{18}F ; or wherein R^4 is ^{123}I , ^{125}I , ^{75}Br , ^{76}Br , ^{77}Br or ^{18}F . More preferred are those compounds of formula (I) wherein R^4 is ^{125}I .

10 Another interesting form of radiolabelling a compound of formula (I) is by substituting a carbon atom by a ^{11}C -atom or the substitution of a hydrogen atom by a tritium atom. Introducing such a ^{11}C -atom is conveniently carried out by *N*-alkylating a compound of formula (I), wherein R^1 , R^2 and/or R^3 are hydrogen using a ^{11}C -labelled alkylating reagent. Tritium radiolabelled compounds of formula (I-a) can be prepared by
15 reductively *N*-alkylation intermediates of formula (III) with intermediates of formula (IV) in the presence of $^3\text{H}_2$ -gas as reducing agent and in combination with a suitable catalyst. Those compounds of formula (I) containing a halogen atom may also be suitably converted to tritium radiolabelled compounds of formula (I) by exchanging the halogen atom with $^3\text{H}_2$ -gas in the presence of a suitable catalyst such as, for example
20 palladium on activated carbon.

Hence, said radiolabelled compounds of formula (I) can be used in a process of specifically marking dopamine D_4 receptor sites in biological material. Said process comprises the steps of (a) radiolabelling a compound of formula (I), (b) administering this radiolabelled compound to biological material and subsequently (c) detecting the
25 emissions from the radiolabelled compound. The term biological material is meant to comprise every kind of material which has a biological origin. More in particular this term refers to tissue samples, plasma or body fluids but also to animals, specially warm-blooded animals, or parts of animals such as organs.

The radiolabelled compounds of formula (I) are also useful as agents for screening
30 whether a test compound has the ability to occupy or bind to a dopamine D_4 receptor site. The degree to which a test compound will displace a compound of formula (I) from the dopamine D_4 receptor site will show the test compound ability as either an agonist, an antagonist or a mixed agonist/antagonist of a dopamine D_4 receptor.

When used in *in vivo* assays, the radiolabelled compounds are administered in an
35 appropriate composition to an animal and the location of said radiolabelled compounds is detected using imaging techniques, such as, for instance, Single Photon Emission Computerized Tomography (SPECT) or Positron Emission Tomography (PET) and the

-14-

like. In this manner the distribution of dopamine D₄ receptor sites throughout the body can be detected and organs containing dopamine D₄ receptor sites such as, for example, the brain, can be visualized by the imaging techniques mentioned hereinabove. This process of imaging an organ by administering a radiolabelled compound of formula (I), which binds to the dopamine D₄ receptor sites and detecting the emissions from the radioactive compound also constitutes a part of the present invention.

The following examples are intended to illustrate and not to limit the scope of the present invention.

10 A. Preparation of the intermediates

Example A.1

a) A mixture of ethyl 4-(methylamino)-1-piperidinecarboxylate (111.75 g) and 2-chloro-*N,N*-dimethyl-4-pyrimidinamine (47.3 g) is stirred and heated in an oil-bath for 22 hours at 120°C. The reaction mixture is cooled and the product is taken up in trichloromethane (500 ml). Water (300 ml) is added and the layers are separated. The organic phase is washed with water (200 ml), dried, filtered and evaporated. The residue solidifies on triturating in petroleumether. The product is filtered off and crystallized from diisopropyl ether (390 ml). After cooling to 0°C, the product is filtered off and dried, yielding 55.5 g (59.7%) of ethyl 4-[[4-dimethylamino-2-pyrimidinyl]methylamino]-1-piperidinecarboxylate (intermediate 1).

b) A mixture of intermediate 1 (52.5 g), potassium hydroxide (95.4 g) and 2-propanol (950 ml) is stirred and refluxed for 18 hours. The reaction mixture is evaporated and water (900 ml) are added to the residue. The whole is stirred for 30 minutes in a boiling water-bath and evaporation is continued till all traces of 2-propanol are removed. After cooling, the product is extracted twice with dichloromethane (270 ml). The combined extracts are washed with 280 parts of diisopropyl ether (390 ml). After cooling to 0°C, the product is filtered off and dried, yielding 29 g (72.5%) of *N*²,*N*⁴,*N*⁴-trimethyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (intermediate 2)

In a similar way were prepared :

*N*⁴-ethyl-*N*²-methyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 3);
*N*⁴-cyclopropyl-*N*²-methyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 4);
N-methyl-*N*-(4-piperidinyl)-4-(1-pyrrolidinyl)-2-pyrimidinamine (interm. 5);
*N*²,*N*⁴-dimethyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 6);
*N*²-butyl-*N*⁴,*N*⁴-dimethyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 7);
*N*⁴,*N*⁴-dimethyl-*N*²-methyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 8);
*N*²-methyl-*N*²-(4-piperidinyl)-*N*⁴,*N*⁴-dipropyl-2,4-pyrimidinediamine (interm. 9);
*N*²-ethyl-*N*⁴,*N*⁴-dimethyl-*N*²-(4-piperidinyl)-2,4-pyrimidinediamine (interm. 10);

-15-

*N*⁴,*N*⁴-dimethyl-*N*²-(4-piperidiny)-2,4-pyrimidinediamine (interm. 11);
*N*²-methyl-*N*²-(4-piperidiny)-2,4-pyrimidinediamine (interm. 12);
*N*⁴-butyl-*N*²-methyl-*N*²-(4-piperidiny)-2,4-pyrimidinediamine (interm. 13); and
*N*²-methyl-*N*⁴-(1-methylethyl)-*N*²-(4-piperidiny)-2,4-pyrimidinediamine (interm. 14).

Example A.2

A solution of intermediate (7) (6.8 g) in 2-propanol (32 ml) was acidified with concentrated nitric acid. The crystallized product was filtered off and dried, yielding 7 g (70.7%) of *N*²-butyl-*N*⁴,*N*⁴-dimethyl-*N*²-(4-piperidiny)-2,4-pyrimidinediamine dinitrate (interm. 15; mp. 168.6°C).

In a similar way, the dinitrate salt of intermediates (8) and (9) were prepared.

B. Preparation of the final compounds

Example B.1

A mixture of bromomethylbenzene (0.06 g), intermediate 3 (0.100 g) and sodium carbonate (0.100 g) in 4-methyl-2-pentanone (2 ml) was stirred overnight at 60°C. The mixture was filtered, diluted with CH₂Cl₂ until a 15 ml total volume was obtained, then purified by high-performance liquid chromatography over silica gel (eluent : A = CH₂Cl₂ and B = CH₂Cl₂/CH₃OH 90/10 : from 100% A to 100% B over 20 minutes; 125 ml/min). The desired fractions were collected and the solvent was evaporated, yielding 0.03 g of *N*⁴-ethyl-*N*²-methyl-*N*²-[1-(phenylmethyl)-4-piperidiny]-2,4-pyrimidinediamine (comp. 6)

Example B.2

Using the same procedure as described in example B.1, but replacing 4-methyl-2-pentanone by *N,N*-dimethyl-formamide, intermediate 2 was reacted with 1-(2-chloroethyl)-3-methoxybenzene, to form *N*²-[1-[2-(3-methoxyphenoxy)ethyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidinediamine (comp. 26, mp. 104.8°C).

Example B.3

Using the same procedure as describe in example B.1 and adding a small amount of potassium iodide to the reaction mixture, intermediate 2 was reacted with 1,1'-(4-chlorobutylidene)bis[4-fluorobenzene] to form *N*²-[1-[4,4-bis(4-fluorophenyl)butyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidinediamine (comp. 44, mp. 106.3°C).

Example B.4

A mixture of chloromethylbenzene (1.9 g), intermediate 2 (3 g) and triethylamine

-16-

(2.62 g) in *N,N*-dimethylacetamide (55 ml) was stirred for 2 hours at 75°C. The reaction mixture was poured out into ice-water and extracted with toluene. The separated organic layer was washed three times with water, dried, filtered and the solvent evaporated. The residue was dissolved in boiling petroleum ether and treated with activated charcoal. The mixture was filtered warm over dicalite, and the filtrate was concentrated. The concentrate was cooled on an ice-bath and the resulting precipitate was filtered off and dried, yielding 2.3 g (54.4%) of *N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-(phenylmethyl)-4-piperidinyl]-2,4-pyrimidinediamine (comp. 8).

10 Example B.5

A mixture of intermediate 4 (4.9 g) and 1,3-benzodioxole-5-carboxaldehyde (3 g) in methanol (250 ml) was hydrogenated at 50°C with platinum on activated carbon (2 g) as a catalyst in the presence of thiophene (4%; 1 ml). After uptake of H₂, the catalyst was filtered off over dicalite, and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 90/10). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in methanol and converted into the ethanedioic acid salt (1:2). The precipitate was filtered off and dried, yielding 7.43 g (66.2%) of *N*²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidinyl]-*N*⁴-cyclopropyl-*N*²-methyl-2,4-pyrimidinediamine ethanedioate(1:2) (comp. 67).

Example B.6

A mixture of 1-(2-phenylethyl)-4-piperidinamine (8.17 g) and 2-chloro-*N,N*-dimethyl-4-pyrimidinamine (6.3 g) was stirred for 20 hours at 120°C. The reaction mixture was diluted with water and extracted with CH₂Cl₂. The organic layer was washed with water, dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH(NH₃) 96/4). The pure fractions were collected and the solvent evaporated. The residue was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding 3.69 g (28.4%) of *N*⁴,*N*⁴-dimethyl-*N*²-[1-(2-phenylethyl)-4-piperidinyl]-2,4-pyrimidinediamine (comp. 11).

Example B.7

A mixture of compound 67 (0.005 mol) and calcium carbonate (0.0065 mol) in CH₂Cl₂ (25 ml) and methanol (10 ml) was stirred at room temperature. A solution of benzyltrimethylammonium dichloroiodate (0.005 mol) in CH₂Cl₂ (5 ml) and methanol (5 ml) was added dropwise at room temperature. The reaction mixture was stirred at room temperature for 30 minutes, then washed with water. The organic layer was separated, dried, filtered, and the solvent was evaporated. The residue was purified by


-17-

column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue (1.6 g) was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (3:2) with (E)-2-butenedioic acid. The precipitate was filtered off and dried, yielding 1.28 g (41.1%) of N²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-N⁴-cyclopropyl-N²-methyl-5-iodo-2,4-pyrimidinediamine . (E)-2-butenedioate (2:3) hydrate (1:1) (comp. 88).

Example B.8

A mixture of compound 67 (1.6 mg), acetic acid (0.5 ml; 100%), I¹²⁵ (100 μCu) and H₂O₂ (0.1 ml) was stirred at room temperature. The reaction was quenched with sodiumsulfite solution (1.8 ml; 1M) , yielding N²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-N⁴-cyclopropyl-N²-methyl-5-iodo-¹²⁵I-2,4-pyrimidinediamine (comp. 89).

Tables 1 to 5 list compounds of formula (I) which were prepared according to one of the above examples.

15 Table 1



Co. No.	Ex No.	R ^x	R ^y	Alk	R ¹	R ²	R ³	Phys. Data
1	B.1	H	H	(CH ₂) ₄	CH ₃	CH ₃	CH ₃	-
2	B.1	H	H	(CH ₂) ₃	CH ₃	CH ₃	CH ₃	-
3	B.1	H	H	CH(CH ₃) ₂	CH ₃	CH ₃	CH ₃	-
4	B.1	H	H	CH ₂	(CH ₂) ₃ CH ₃	CH ₃	CH ₃	-
5	B.1	H	H	CH ₂	CH ₃	cyclopropyl	H	-
6	B.1	H	H	CH ₂	CH ₃	CH ₂ CH ₃	H	-
7	B.3	H	H	CH=CHCH ₂	CH ₃	CH ₃	CH ₃	(E)-2-butenedioate / H ₂ O
8	B.4	H	H	CH ₂	CH ₃	CH ₃	CH ₃	-
9	B.5	H	H	CH ₂	CH ₃	CH ₂ CH ₃	CH ₂ CH ₃	-
10	B.5	H	H	CH ₂	CH ₃	CH ₃	H	-
11	B.6	H	H	(CH ₂) ₂	H	CH ₃	CH ₃	-
12	B.5	H	3-CH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
13	B.1	H	4-phenyl	CH ₂	CH ₃	CH ₃	CH ₃	-
14	B.1	H	2-Cl	CH ₂	CH ₃	CH ₃	CH ₃	-

-18-

Co. No.	Ex No.	R ^x	R ^y	Alk	R ¹	R ²	R ³	Phys. Data
16	B.1	H	2-F	CH ₂	CH ₃	CH ₃	CH ₃	-
17	B.1	H	3-F	CH ₂	CH ₃	CH ₃	CH ₃	-
18	B.1	H	3-CF ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
19	B.1	H	4-NO ₂	CH ₂	CH ₃	CH ₃	CH ₃	-
20	B.1	H	4-Cl	CH ₂	CH ₃	CH ₃	CH ₃	-
21	B.1	H	3-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
22	B.1	H	4-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
23	B.1	H	3-Cl	CH ₂	CH ₃	CH ₃	CH ₃	-
24	B.1	H	4-F	CH ₂	CH ₃	CH ₃	CH ₃	-
25	B.2	H	2-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	mp. 168.4°C/ .2(COOH) ₂
26	B.2	H	3-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	mp. 104.8°C
27	B.4	H	4-Cl	CH ₂	CH ₃	CH ₃	CH ₃	-
28	B.5	H	2-CH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
29	B.5	H	4-CH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
30	B.5	H	2-CF ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
31	B.5	H	4-CF ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
32	B.5	H	2-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
33	B.5	H	4-OH	CH ₂	CH ₃	CH ₃	CH ₃	-
34	B.5	H	4-CH(CH ₃) ₂	CH ₂	CH ₃	CH ₃	CH ₃	-
35	B.1	2-CH ₃	5-CH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
36	B.1	2-Cl	6-Cl	CH ₂	CH ₃	CH ₃	CH ₃	-
37	B.5	3-OCH ₃	4-OCH ₃	CH ₂	CH ₃	CH ₃	CH ₃	-
38	B.5	H	H	CH ₂	CH ₃	-(CH ₂) ₄ -*		

* : R² and R³ are taken together to form a bivalent radical


Table 2



Co. No.	Ex. No.	R ¹	R ²	R ³	Physical data
39	B.3	CH ₃	H	H	mp. 253.4°C/.2 (COOH) ₂
40	B.3	CH ₃	H	CH ₃	mp. 111.8°C
41	B.3	CH ₃	cyclopropyl	H	mp. 168.9°C/.2 (COOH) ₂
42	B.3	CH ₃	CH ₂ CH ₃	H	mp. 168.9°C/.2 (COOH) ₂
43	B.3	CH ₃	(CH ₂) ₃ CH ₃	H	mp. 186.7°C/.2 (COOH) ₂
44	B.3	CH ₃	CH ₃	CH ₃	mp. 106.3°C
45	B.3	CH ₃	CH(CH ₃) ₂	H	mp. 177.1°C/.3/2 (COOH) ₂
46	B.3	(CH ₂) ₃ CH ₃	CH ₃	CH ₃	mp. 102.1°C
47	B.3	CH ₃	-CH ₂ -CH ₂ -CH ₂ -*		mp. 173.2°C/.2 (COOH) ₂

* : R² and R³ are taken together to form a bivalent radical

Table 3




Co. No.	Ex. No.	Alk	R ¹	R ²	R ³	Physical data
48	B.1	CH ₂	H	CH ₃	CH ₃	mp. 126.9°C
49	B.4	CH ₂	CH ₃	H	H	mp. 147.6°C
50	B.1	CH ₂	CH ₃	CH ₃	H	mp. 128.8°C
51	B.4	CH ₂	CH ₃	CH ₃	CH ₃	mp. 97.4°C
52	B.1	CH ₂	CH ₃	(CH ₂) ₃ CH ₃	H	mp. 189.1°C/2(COOH) ₂
53	B.1	CH ₂	CH ₃	CH(CH ₃) ₂	H	mp. 194.2°C/2(COOH) ₂
54	B.1	CH ₂	CH ₃	CH ₂ CH ₃	H	mp. 170°C/3/2(COOH) ₂
55	B.1	CH ₂	CH ₃	cyclopropyl	H	mp. 166°C/ .3/2 (COOH) ₂ .H ₂ O
56	B.1	CH ₂	CH ₃	CH ₂ CH ₃	CH ₂ CH ₃	mp. 153.9°C/ .2 (COOH) ₂ .H ₂ O
57	B.1	CH ₂	CH ₂ CH ₃	CH ₃	CH ₃	mp. 141.7°C/ .2 (COOH) ₂ .H ₂ O
58	B.1	CH ₂	(CH ₂) ₃ CH ₃	CH ₃	CH ₃	mp. 132.5°C/2(COOH) ₂
59	B.1	CH ₂	(CH ₂) ₂ CH ₃	(CH ₂) ₂ CH ₃	H	mp. 132.8°C/2(COOH) ₂
60	B.5	CH(CH ₃)	CH ₃	CH ₃	CH ₃	mp. 211°C/ .2(E)-2-butenedioate
61	B.1	CH ₂	CH ₃	-CH ₂ -CH ₂ -CH ₂ -*		mp. 107°C

* : R² and R³ are taken together to form a bivalent radical

-20-


Table 4





Co. No.	Ex. No.	R ¹	R ²	R ³	Physical data
62	B.5	CH ₃	CH ₃	H	-
63	B.4	CH ₃	CH ₃	CH ₃	mp. 127.1°C
64	B.5	CH ₃	(CH ₂) ₂ CH ₃	(CH ₂) ₂ CH ₃	-
65	B.5	CH ₃	CH ₂ CH ₃	CH ₂ CH ₃	-
66	B.5	CH ₃	cyclopropyl	H	-
67	B.5	CH ₃	cyclopropyl	H	ethandioate (1:2)
68	B.5	CH ₃	CH ₂ CH ₃	H	-
69	B.5	(CH ₂) ₃ CH ₃	CH ₃	CH ₃	-
70	B.5	CH ₃	-CH ₂ -CH ₂ -CH ₂ -CH ₂ -*		-

* : R² and R³ are taken together to form a bivalent radical

5 Table 5



Co. No.	Ex. No.	Q	Alk	Physical data
71	B.3	phenoxy	-CH(CH ₃)CH ₂ -	.2 (COOH) ₂ .H ₂ O ; mp. 142.5°C
72	B.4	phenoxy	(CH ₂) ₂	.2 (COOH) ₂ ; mp. 192.6°C
73	B.4	phenoxy	(CH ₂) ₄	.2 (COOH) ₂ ; mp. 162.8°C
74	B.4	4-fluorophenoxy	(CH ₂) ₃	mp. 113.8°C
75	B.4	2-acetylphenoxy	(CH ₂) ₂	mp. 133.2°C
76	B.4	2-chloro-6-methylphenoxy	(CH ₂) ₂	.2 (COOH) ₂ ; mp. 171.9°C
77	B.1	2-pyridinyl	CH ₂	-
78	B.5	3-pyridinyl	CH ₂	-
79	B.5	4-pyridinyl	CH ₂	-
80	B.5	2-thienyl	CH ₂	-
81	B.1	1-naphthalenyl	CH ₂	-

Co. No.	Ex. No.	Q	Alk	Physical data
83	B.5	2-naphthalenyl	CH ₂	-
84	B.1	2-quinoliny	CH ₂	-
85	B.4	3-indolyl	(CH ₂) ₂	mp. 201.2°C
86	B.4		CH ₂	mp. 142.1°C
87	B.2		CH ₂	.3/2 (E)-2-butenedioate ; mp. 226.3°C

C. Pharmacological example

Example C.1. : *In vitro* binding affinity for dopamine D₄ receptor

The interaction of the compounds of formula (I) with the dopamine D₄ receptors was assessed in *in vitro* radioligand binding experiments. A low concentration of ³H-spiperone with a high binding affinity for the dopamine D₄ receptor was incubated with a sample of a membrane preparation of transfected Chinese Hamster Ovary (CHO) cells which express cloned human D₄ receptors (Receptor Biology, Maryland, USA) in a buffered medium. When equilibrium of binding was reached, the receptor bound radioactivity was separated from the non-bound radioactivity, and the receptor-bound activity was counted. The interaction of the test compounds, added to the incubation mixture in various concentrations, with the dopamine D₄ receptor was assessed in competition binding experiments as described by Schotte *et al.* (Psychopharmacology, 1996). The compounds with number 1, 2, 5-10, 12-14, 16-24, 26-32, 34-45, 52, 53, 55, 59, 61-63, 66, 68, 70, 73-76, 78, 80, 81 and 83-88 had a pIC₅₀ greater than or equal to 7 (pIC₅₀ is defined as -log IC₅₀ wherein IC₅₀ is the concentration of the test compound causing an inhibition of 50 % of the dopamine D₄ receptors).

D. Composition examples

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.

Example D.1 : Capsules

20 g of the A.I., 6 g sodium lauryl sulfate, 56 g starch, 56 g lactose, 0.8 g colloidal silicon dioxide, and 1.2 g magnesium stearate are vigorously stirred together. The resulting mixture is subsequently filled into 1000 suitable hardened gelatin capsules,

-22-

each comprising 20 mg of the A.I..

Example D.2 : Film-coated tablets

Preparation of tablet core

- 5 A mixture of 100 g of the A.I., 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there are added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole is mixed well and compressed into tablets,
10 giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

- To a solution of 10 g methyl cellulose in 75 ml of denaturated ethanol there is added a solution of 5 g of ethyl cellulose in 150 ml of dichloromethane. Then there are added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol is
15 molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the former and then there are added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated colour suspension and the whole is homogenated. The tablet cores are coated with the thus obtained mixture in a coating apparatus.

20 Example D.3 : Oral solution


- 9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 l of boiling purified water. In 3 l of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 l
25 1,2,3-propanetriol and 3 l of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable
30 containers.

Example D.4 : Injectable solution

- 1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50°C there
35 were added while stirring 4 grams lactic acid, 0.05 grams propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 l, giving a solution comprising 4 mg/ml of A.I.. The solution was sterilized by filtration and filled in sterile containers.

Claims

1. A compound having the formula




- 5 a *N*-oxide form, a pharmaceutically acceptable acid addition salt or a stereochemically isomeric form thereof, wherein
- Alk is C₁₋₆alkanediyl or C₃₋₆alkenediyl;
- R¹ is hydrogen or C₁₋₄alkyl;
- R² and R³ each independently are hydrogen, C₁₋₆alkyl or C₃₋₇cycloalkyl; or
- 10 R² and R³ may also be taken together with the nitrogen atom to which they are attached, thus forming a pyrrolidine, a piperidine or a perhydro azepine ring;
- R⁴ is hydrogen or halo;
- Q is aryl, aryloxy, di(aryl)methyl or heteroaryl;
- aryl is naphthyl or phenyl, said naphthyl and phenyl may optionally be substituted
- 15 with one, two or three substituents selected from halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkylcarbonyl, haloC₁₋₄alkyl, nitro, amino, cyano and phenyl; and
- heteroaryl is quinolinyl, isoquinolinyl, pyridinyl, thienyl, indolyl, 2,3-dihydro-1,4-benzodioxinyl, 2,3-dihydro-benzofuranyl or benzodioxolanyl; said heteroaryls
- 20 may optionally be substituted with one, two or three substituents selected from halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkylcarbonyl, haloC₁₋₄alkyl and phenyl.
2. A compound according to claim 1 wherein R⁴ is hydrogen.
- 25 3. A compound according to claim 1 or 2 wherein Alk is a straight chained C₁₋₆alkanediyl and Q is optionally substituted phenyl, 2-naphthyl, 2-quinolinyl, 2,3-dihydrobenzofuran-5-yl or 5-benzodioxolanyl.
4. A compound according to any one of claims 1 to 3 wherein R¹ is C₁₋₄alkyl; R² is C₁₋₆alkyl or C₃₋₇cycloalkyl and R³ is hydrogen, C₁₋₆alkyl; or R² and R³ form a
- 30 pyrrolidine ring when taken together with the nitrogen atom to which they are attached.
5. A compound according to claim 1 wherein the compounds is :

-24-


- N*⁴-cyclopropyl-*N*²-methyl-*N*²-[1-(phenylmethyl)-4-piperidiny]-2,4-pyrimidine-diamine;
- N*²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-*N*⁴-ethyl-*N*²-methyl-2,4-pyrimidinediamine;
- 5 *N*²-[1-(1,3-benzodioxol-5-ylmethyl)-4-piperidiny]-*N*⁴-cyclopropyl-*N*²-methyl-2,4-pyrimidinediamine;
- N*²-[1-[(4-chlorophenyl)methyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidine diamine;
- N*²-[1-[(3-fluorophenyl)methyl]-4-piperidiny]-*N*²,*N*⁴,*N*⁴-trimethyl-2,4-pyrimidine-
10 diamine;
- N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-[[4-(trifluoromethyl)phenyl]methyl]-4-piperidiny]-2,4-pyrimidinediamine;
- N*²,*N*⁴,*N*⁴-trimethyl-*N*²-[1-[(4-methylphenyl)methyl]-4-piperidiny]-2,4-pyrimidine-
15 diamine; a stereochemically isomeric form or a pharmaceutically acceptable addition salt thereof.
6. A compound according to claim 1 having at least one halo which is a radioactive
20 isotope of iodine, bromine or fluorine, or having at least one ¹¹C-atom or tritium atom.
7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier
and, as active ingredient, an effective dopamine D₄ receptor antagonizing amount of
25 a compound as claimed in any one of the claims 1 to 6.
8. A process of preparing a pharmaceutical composition as defined in claim 7,
characterized in that, a pharmaceutically acceptable carrier is intimately mixed with
an effective dopamine D₄ receptor antagonizing amount of a compound as claimed
30 in any one of claims 1 to 6.
9. A compound as claimed in any one of claims 1 to 5 for use as a medicine.
10. Use of a compound as claimed in any one of claims 1 to 5 for the manufacture of a
35 medicament for treating psychotic disorders.
11. A process of preparing a compound as claimed in claim 1, characterized by
a) *N*-alkylating an intermediate of formula

-25-




wherein R^1 , R^2 , R^3 and R^4 are defined as in claim 1, with an intermediate of formula Q-Alk-W (II) wherein W represents an appropriate reactive leaving group and Alk is defined as in claim 1, in a reaction-inert solvent and in the presence of a suitable base,
 5 and optionally in the presence of potassium iodide;

b) reacting an intermediate of formula (V)




wherein Q, Alk and R^1 are defined as in claim 1, with an intermediate of formula (VI)




10 wherein R^2 , R^3 and R^4 are defined as in claim 1 and W^1 is a suitable leaving group;

c) reductively N-alkylating an intermediate of formula (III)



wherein R^1 , R^2 , R^3 and R^4 are defined as in claim 1, with an aldehyde or ketone of formula Q-Alk'=O (IV), said Q-Alk'=O being derived from Q-Alk'H₂ by replacing two geminal hydrogen atoms by an oxo group in a reaction-inert solvent and in the presence
 15 of a reducing agent; thus forming a compound of formula (I-a);



wherein Alk'H is the same as Alk defined as in claim 1 but whereby Alk is attached to the nitrogen of the piperidine ring by a carbon atom bearing at least one hydrogen
 20 atom;

d) converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or conversely, converting the acid addition salt form into the free base by treatment with alkali; and, if desired, preparing stereochemically isomeric forms and/or *N*-oxides thereof.

12. A process of marking a dopamine D₄ receptor comprising the steps of
 - a) radiolabelling a compound as claimed in claim 1;
 - b) administering said radiolabelled compound to biological material,
 - c) detecting the emissions from the radiolabelled compound.
13. A process of imaging an organ, characterized by, administering a sufficient amount of a radiolabelled compound of formula (I) in an appropriate composition, whereby said compound binds to the dopamine D₄ receptor sites; and detecting the emissions from the radioactive compound.

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.
PCT/EP 97/02506

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D401/12 C07D405/14 C07D401/14 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 10018 A (NEUROGEN CORP ;THURKAUF ANDREW (US); HUTCHISON ALAN (US)) 4 April 1996 see page 19 - page 20 ---	1-13
A	EP 0 532 178 A (ICI PLC) 17 March 1993 see page 2, line 1 - line 6; example 91 ---	1-13
A	WO 93 17017 A (JANSSEN PHARMACEUTICA NV) 2 September 1993 cited in the application see claim 1; examples 125-128,132; tables 3,4 --- -/--	1-13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

2 September 1997

Date of mailing of the international search report

15.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/02506

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 379 806 A (MITSUI PETROCHEMICAL IND ;MITSUI PHARMACEUTICALS (JP)) 1 August 1990 cited in the application see page 29, no 2198 -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/02506

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9610018 A	04-04-96	US 5633376 A	27-05-97
		AU 3725995 A	19-04-96
		EP 0783494 A	16-07-97
		NO 971423 A	23-05-97
		US 5646279 A	08-07-97
		US 5646280 A	08-07-97
		AU 4368996 A	17-06-96
		WO 9616040 A	30-05-96

EP 0532178 A	17-03-93	AT 144254 T	15-11-96
		AU 645707 B	20-01-94
		AU 2101592 A	18-02-93
		CA 2076146 A	16-02-93
		DE 69214587 D	21-11-96
		DE 69214587 T	20-03-97
		HU 66515 A	28-12-94
		JP 5239026 A	17-09-93
		US 5455246 A	03-10-95
		US 5266570 A	30-11-93

WO 9317017 A	02-09-93	AP 416 A	29-09-95
		AT 138064 T	15-06-96
		AU 3499193 A	13-09-93
		BG 98870 A	31-03-95
		CA 2117483 A	02-09-93
		CN 1079470 A	15-12-93
		CZ 9402020 A	18-01-95
		DE 69302687 D	20-06-96
		DE 69302687 T	14-11-96
		EP 0639192 A	22-02-95
		ES 2087721 T	16-07-96
		FI 943928 A	26-08-94
		HU 71129 A	28-11-95
		HU 9500317 A	28-09-95
		JP 7504408 T	18-05-95
		LT 3049 B	25-10-94
		LV 10715 B	20-12-95
		NO 943186 A	29-08-94
		NZ 249124 A	28-08-95
		SK 102994 A	11-07-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/02506

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9317017 A		US 5541180 A	30-07-96
		US 5607949 A	04-03-97
		ZA 9301404 A	26-08-94

EP 0379806 A	01-08-90	JP 2221275 A	04-09-90
		AT 136542 T	15-04-96
		AU 629595 B	08-10-92
		AU 4732989 A	05-07-90
		CA 2006944 A	29-06-90
		DE 68926232 D	15-05-96
		DE 68926232 T	17-10-96
		EP 0612746 A	31-08-94
		HU 210001 B	30-01-95
		US 5147876 A	15-09-92
		US 5264435 A	23-11-93
		CN 1045390 A	19-09-90
		CN 1090846 A	17-08-94
		HU 209594 B	29-08-94
		HU 209574 B	29-08-94
		JP 3014568 A	23-01-91

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.